

Literaturverzeichnis

Hamburger Ärzteblatt 05 | 2019

Seite 1

Weidestr. 122 b
22083 Hamburg
Redaktion
E-Mail: verlag@aeckhh.de
Tel.: (040) 20 22 99 – 205
Fax: (040) 20 22 99 – 400

S. 12–17: Multiple Sklerose: Verlauf und Stadium bestimmen die Therapie

Von Dr. Thorsten Rosenkranz, PD Dr. Christoph Terborg

1. Runmarker B, Andersson O. Prognostic factors in a multiple sclerosis incidence cohort with 25 years follow-up. *Brain*. 1993;116:117–34.
2. Thompson AJ, Banwell BL, Barkhof F, et al. Diagnosis of multiple sclerosis: 2017 revisions of the McDonald criteria. *Lancet Neurol*. 2018;17:162–173.
3. Burton JM, O'Connor PW, Hohol M, et al. Oral versus intravenous steroids for treatment of relapses in multiple sclerosis. *Cochrane Database Syst Rev*. 2012;12 CD006921.
4. Scolding NJ, Pasquini M, Reingold SC, et al. Cell based therapeutic strategies for multiple sclerosis. *Brain*. 2017; 140:2776–2796.
5. Willekens B und Cools N. Beyond the magic bullet: Current progress of therapeutic vaccination in multiple sclerosis. *CNS Drugs*. 2018; 32:401–410.

S. 30–33: Wenn die Leiste „zwickt“ und es doch kein Leistenbruch ist ...

Von Dr. Christian Gatzka, PD Dr. Christian Tesch, PD Dr. Lorenz Büchler

1. Ganz R, Parvizi J, Beck M, Leunig M, Nötzli H, Siebenrock KA. Femoroacetabular impingement: a cause for osteoarthritis of the hip. *Clin Orthop Relat Res*. 2003 Dec;417:112–20.
2. Albers CE, Steppacher SD, Haefeli PC, Werlen S, Hanke MS, Siebenrock KA, et al. Twelve percent of hips with a primary cam deformity exhibit a slip-like morphology resembling sequelae of slipped capital femoral epiphysis. *Clin Orthop Relat Res*. 2015 Apr;473(4):1212–23.
3. van Klij P, Heerey J, Waarsing JH, Agricola R. The Prevalence of Cam and Pincer Morphology and Its Association With Development of Hip Osteoarthritis. *J Orthop Sports Phys Ther*. 2018 Apr;48(4):230–8.
4. Kang AC, Gooding AJ, Coates MH, Goh TD, Armour P, Rietveld J. Computed tomography assessment of hip joints in asymptomatic individuals in relation to femoroacetabular impingement. *Am J Sports Med*. 2010 Jun;38(6):1160–5.
5. Agricola R, Waarsing JH, Arden NK, Carr AJ, Bierma-Zeinstra SM, Thomas GE, et al. Cam impingement of the hip: a risk factor for hip osteoarthritis. *Nat Rev Rheumatol*. 2013 Oct;9(10):630–4.
6. Büchler L, Kindler M, Dienst M, Lampert C, Fickert S, Möckel G, et al. Diagnostik des Hüftgelenkes. AGA, editor. Zürich: AGA - Hüftkomitee; 2017.
7. Nötzli HP, Wyss TF, Stoecklin CH, Schmid MR, Treiber K, Hodler J. The contour of the femoral head-neck junction as a predictor for the risk of anterior impingement. *J Bone Joint Surg Br*. 2002 May;84(4):556–60.
8. Hartofilakidis G, Bardakos NV, Babis GC, Georgiades G. An examination of the association between different morphotypes of femoroacetabular impingement in asymptomatic subjects and the development of osteoarthritis of the hip. *J Bone Joint Surg Br*. 2011 May;93(5):580–6.
9. Philippon MJ, Briggs KK, Yen YM, Kuppersmith DA. Outcomes following hip arthroscopy for femoroacetabular impingement with associated chondrolabral dysfunction: minimum two-year follow-up. *J Bone Joint Surg Br*. 2009 Jan;91(1):16–23.
10. Siebenrock KA, Ferner F, Noble PC, Santore RF, Werlen S, Mamisch TC. The cam-type deformity of the proximal femur arises in childhood in response to vigorous sporting activity. *Clin Orthop Relat Res*. 2011 Nov;469(11):3229–40.
11. Lee WY, Kang C, Hwang DS, Jeon JH, Zheng L. Descriptive Epidemiology of Symptomatic Femoroacetabular Impingement in Young Athlete: Single Center Study. *Hip Pelvis*. 2016 Mar;28(1):29–34.

Literaturverzeichnis

Hamburger Ärzteblatt 05 | 2019

Seite 2

Weidestr. 122 b
22083 Hamburg
Redaktion
E-Mail: verlag@aekhh.de
Tel.: (040) 20 22 99 – 205
Fax: (040) 20 22 99 – 400

12. Hanke MS, Steppacher SD, Zurmuhle CA, Siebenrock KA, Tannast M. Hips With Protrusio Azetabuli Are at Increased Risk for Failure After Femoroazetabular Impingement Surgery: A 10-year Followup. *Clin Orthop Relat Res.* 2016 Oct;474(10):2168-80.
13. Albers CE, Steppacher SD, Ganz R, Tannast M, Siebenrock KA. Impingement adversely affects 10-year survivorship after periazetabular osteotomy for DDH. *Clin Orthop Relat Res.* 2013 May;471(5):1602-14.
14. Tönnis D, Heinecke A. Azetabular and femoral anteversion: relationship with osteoarthritis of the hip. *J Bone Joint Surg Am.* 1999 Dec;81(12):1747-70.
15. Clohisy JC, St John LC, Nunley RM, Schutz AL, Schoenecker PL. Combined periazetabular and femoral osteotomies for severe hip deformities. *Clin Orthop Relat Res.* 2009 Sep;467(9):2221-7.
16. Maier C, Zingg P, Seifert B, Sutter R, Dora C. Femoral torsion: reliability and validity of the trochanteric prominence angle test. *Hip Int.* 2012 Sep-Oct;22(5):534-8.
17. Domb BG, Nasser RM, Botser IB. Partial-thickness tears of the gluteus medius: rationale and technique for trans-tendinous endoscopic repair. *Arthroscopy.* 2010 Dec;26(12):1697-705.
18. Tannast M, Siebenrock KA, Anderson SE. Femoroazetabular impingement: radiographic diagnosis--what the radiologist should know. *AJR Am J Roentgenol.* 2007 Jun;188(6):1540-52.
19. Schmaranzer F, Klauser A, Kogler M, Henninger B, Forstner T, Reichkendler M, et al. MR arthrography of the hip with and without leg traction: Assessing the diagnostic performance in detection of ligamentum teres lesions with arthroscopic correlation. *Eur J Radiol.* 2016 Feb;85(2):489-97.
20. Botser IB, Ozoude GC, Martin DE, Siddiqi AJ, Kuppuswami S, Domb BG. Femoral anteversion in the hip: comparison of measurement by computed tomography, magnetic resonance imaging, and physical examination. *Arthroscopy.* 2012 May;28(5):619-27.
21. Tesch C. Sonografische Untersuchung des Labrum azetabulare beim Hüftschmerz -Sonografische Untersuchungstechnik-. OUP. 2015;4(3):3.
22. Weinreb JH, Sheth C, Apostolakos J, McCarthy MB, Barden B, Cote MP, et al. Tendon structure, disease, and imaging. *Muscles Ligaments Tendons J.* 2014 Jan;4(1):66-73.
23. Mei-Dan O, McConkey MO, Brick M. Catastrophic failure of hip arthroscopy due to iatrogenic instability: can partial division of the ligamentum teres and iliofemoral ligament cause subluxation? *Arthroscopy.* 2012 Mar;28(3):440-5.
24. Harris JD. Hip labral repair: options and outcomes. *Curr Rev Musculoskelet Med.* 2016 Dec;9(4):361-7.
25. Tönnis D, Itoh K, Heinecke A, Behrens K. [The management of congenital hip luxation with arthrographic control, an individual risk-reducing and time-saving method. I. Choice of method and risk assessment based on arthrographic findings]. *Z Orthop Ihre Grenzgeb.* 1984 Jan-Feb;122(1):50-61.
26. Philippon MJ, Schroder ESBG, Briggs KK. Hip arthroscopy for femoroazetabular impingement in patients aged 50 years or older. *Arthroscopy.* 2012 Jan;28(1):59-65.

S. 34 – 35: Arztbesuch hinausgezögert – Diagnose kam zu spät

Von Dr. Anamur Lan Göttinger

1. Buske C, et al. Leitlinie Morbus Waldenström (Lymphoplasmozytisches Lymphom). Deutsche Gesellschaft für Hämatologie und Medizinische Onkologie (DGHO). Stand Dezember 2018
2. Herold G. Innere Medizin 2017. Herold 2016, ISBN 3-981-46606-3.
3. Wörmann B, et al. Leitlinie Multiples Myelom. Deutsche Gesellschaft für Hämatologie und Medizinische Onkologie (DGHO). Stand Mai 2018

Literaturverzeichnis

Hamburger Ärzteblatt 05 | 2019

Seite 3

Weidestr. 122 b
22083 Hamburg
Redaktion
E-Mail: verlag@aeckhh.de
Tel.: (040) 20 22 99 – 205
Fax: (040) 20 22 99 – 400

S. 36–37: Schlimmste Pestseuche seit Menschengedenken

Von Dr. Hans Peter Richter-von Arnault

1. Winkle S. Kulturgeschichte der Seuchen. Pest (422-515), Komet Verlag 1997
2. Bocaccio G. Das Dekameron. Vollständige Ausgabe in der Übersetzung von Karl Witte, bearb. von Helmut Bode, Kunstmuseum Luzern/Freudenstadt 1973
3. Folland CK, Karl TR, KYA Vinnikov. Observed Climate Variation and Change Intergovernmental Panel on Climate Change. In: Houghton JT, Jenkins GJ, Ephraums JJ (Eds). Climate Change: The IPCC. Scientific Assessment, Cambridge: Cambridge University Press, 1990
4. Kehl H. Rekonstruktion der mittleren Temperatur auf der Nordhalbkugel in den vergangenen 1000 bis 2000 Jahren. Vegetationsökologie Tropischer und Subtropischer Klimate. Erläuterungen zur Vorlesung Institut für Ökologie, TU Berlin, 2014