

Literaturverzeichnis

Hamburger Ärzteblatt 12 | 2020

Seite 1

Weidestr. 122 b
22083 Hamburg
Redaktion

E-Mail: verlag@aekhh.de

Tel.: (040) 20 22 99 – 205

Fax: (040) 20 22 99 – 400

S. 12 – 16: 30 Jahre minimalinvasive Bauchchirurgie. *Von Prof. Dr. Carsten Zornig*

1. Semm K. Die endoskopische Appendektomie. Gynäkol Prax 1983;7:26.
2. Perissat J, Collet D, Belliard R. Gallstones. Laparoscopic treatment – cholecystectomy, cholecystotomy and lithotripsy. Surg Endosc 1990;4:1.
3. Götz F, Pier A, Bacher C. Modified laparoscopic appaendectomy in surgey. Surg Endosc 1990;4:6.
4. Zornig C, Emmermann A, von Waldenfels HA, Felixmüller C, Mofid H. Laparoscopic cholecystectomy without visible scar: combined transvaginal and transumbilical approach. Endoscopy 2007;39:913.
5. Nissen R. Eine einfache Operation zur Beeinflussung der Refluxösophagitis. Wochenschrift 1956;86:590.
6. Toupet MA. Technique d'oesophagogastroplastie avec phreno-gastropexie appliquee dans la cure radicale des hernies hiatales et comme complement de l'operation de Heller dans les cardiospasmes. Mem Acad Chir 1963;89:394.
7. Zornig C, Strate U, Fibbe C, Emmermann A, Layer P. Nissen vs. Toupet laparoscopic fundoplication. A prospective randomized study of 200 patients with and without preoperative esophageal motility disorders. Surg Endosc 2002;16.

S. 26 – 28: Hauptstammstenose der linken Herzkranzarterie – wann operieren und wie?

Von PD Dr. Stephan, Dr. Sebastian Bohnen, Prof. Dr. Stephan Willems, Prof. Dr. Berthold Bein und Prof. Dr. Michael Schmoeckel

1. Eckert N, Gießelmann K. Zu wenige Ischämietests. Deutsches Ärzteblatt, Jahrgang 116, 2019; 51-52:1971-73.
2. Fearon WF, Nishi T, De Bruyne B et al. Clinical outcomes and cost-effectiveness of fractional flow reserve-guided percutaneous coronary intervention in patients with stable coronary artery disease: Three-year follow-up of the FAME 2 trial (Fractional Flow Reserve Versus Angiography for Multivessel Evaluation). Circulation 2018;137:480-87.
3. Kirklin JW, Barratt-Boyes B. Cardiac Surgery (Second and Fourth Edition). Churchill Livingstone New York bzw. Elsevier Saunders, Philadelphia; USA (1993 & 2013).
4. Neumann FJ, Sousa-Uva M, Ahlsson A et al. 2018 ESC/EACTS Guidelines on myocardial revascularization. European Heart Journal 2019;40(2):87-165.
5. Stone GW, Kappetein AP, Sabik JF et al. for the EXCEL Trial Investigators. Five-Year Outcomes after PCI or CABG for Left Main Coronary Disease N Engl J Med 2019;381:1820-30.
6. Ricci M, Karamanoukian HL, D'Ancona G et al. Exposure and mechanical stabilization in off-pump coronary artery bypass grafting via sternotomy. Ann Thorac Surg 2000;70(5):1736-40.
7. Beckmann A, Meyer R, Lewandowski J et al. German Heart Surgery Report 2019: The annual updated registry of the German Society for Thoracic and Cardiovascular Surgery. Thorac Cardiovasc Surg 2020;68:263-76.
8. Beckmann A, Meyer R, Lewandowski J et al. German Heart Surgery Report 2017: The annual updated registry of the German Society for Thoracic and Cardiovascular Surgery. Thorac Cardiovasc Surg 2018;66:608-21.
9. Beckmann A, Funkat A, Lewandowski J et al. Cardiac Surgery in Germany during 2012: A report on behalf of the German Society for Thoracic and Cardiovascular Surgery. Thorac Cardiovasc Surg 2014;62:5-17.
10. Dor V, Saab M, Coste P et al. Left ventricular aneurysm: a new surgical approach. Thorac Cardiovasc Surg 1989;37(1),11-19.
11. Geidel S, Laß M, Schneider C et al. Downsizing of the mitral valve and coronary revascularization in severe ischemic mitral regurgitation results in reverse left ventricular and left atrial remodeling. Eur J Cardiothorac Surg 2005;27:1011-16.

Literaturverzeichnis

Hamburger Ärzteblatt 12 | 2020

Seite 2

Weidestr. 122 b

22083 Hamburg

Redaktion

E-Mail: verlag@aekhh.de

Tel.: (040) 20 22 99 – 205

Fax: (040) 20 22 99 – 400

12. Mäkikallio T, Holm NR, Lindsay M et al. On behalf of the NOBLE Study Investigators. Percutaneous coronary angioplasty versus coronary artery bypass grafting in treatment of unprotected left main stenosis (NOBLE): a prospective, randomised, open-label, non-inferiority trial. *Lancet* 2016;388:2743-52.
13. Holm NR, Mäkikallio T, Lindsay MM et al. Percutaneous coronary angioplasty versus coronary artery bypass grafting in the treatment of unprotected left main stenosis: updated 5-year outcomes from the randomised, non-inferiority NOBLE trial. *Lancet* 2020;395:191-9.
14. Thuijs D, Kappetein AP, Serruys PW et al. Percutaneous coronary intervention versus coronary bypass grafting in patients with three-vessel or left main coronary artery disease: 10-year follow-up of the multicenter randomised controlled SYNTAX trial. *Lancet* 2019;394(10206):1325-34.
15. Doenst T, Haverich A, Serruys P et al. PCI and CABG for Treating Stable Coronary Artery Disease. *JACC* 2019;3(8):964-76.
16. Doenst T, Bargenda S, Kirov H et al. Cardiac Surgery 2019 Reviewed *Thorac Cardiovasc Surg* 2020;68:363-76.
17. Diegeler A, Börgermann J, Kappert U et al. Off-pump versus on-pump coronary artery bypass grafting in elderly patients. *N Engl J Med* 2013;368:1189-98.
18. Diegeler A, Börgermann J, Kappert U et al. Five-Year Outcome After Off-Pump or On-Pump Coronary Artery Bypass Grafting in Elderly Patients. *Circulation* 2019;139:1865-71.
19. Lytle BW, Blackstone EH, Sabik JF et al. The effect of bilateral internal thoracic artery grafting on survival during 20 post-operative years. *Ann Thorac Surg* 2004;78(6):2005-12.
20. Iribarne A, Westbrook BM, Malenka DJ et al. Should diabetes be a contraindication to bilateral internal mammary artery grafting? *Ann Thorac Surg* 2018;105(3):709-14.
21. Bayer N. BITA in 9 out of 10 patients - is it safe? Registry results from 2016 to 2020. Lunch Symposium, DGTHG Wiesbaden 02. März 2020.
22. Hassan K, Brüning T, Radke A, Kannmacher J, Bein B, Schmoeckel M, Geidel S. Low Bleeding after emergency CABG using hemadsorption of Ticagrelor: A 2-year clinical experience. *DGTHG-V200; Thoracic Cardiovasc Surg* 2020; 68 (Suppl S1):S47.
23. Hassan K, Kannmacher J, Wohlmuth P, Budde U, Schmoeckel M, Geidel S. Cytosorb adsorption during emergency cardiac operations in patients at high risk of bleeding. *Ann Thoracic Surg* 2019;108:45-51.
24. Hassan K, Stoeck M, Brüning T, Bein B, Caspary M, Schmoeckel M, Geidel S. Budget impact analysis of sorbent hemadsorption during emergency cardiac surgery in Ticagrelor-loaded patients. *DGTHG-KV136; Thoracic Cardiovasc Surg* 2020; 68 (Suppl S1):S67.

S. 30 – 32: Der besondere Fall: Gedächtnislücken wegtherapiert. *Von Joachim W. Simon-Schultz*

1. Blackwell T, Yaffe K, Laffan AM et al. Associations between sleep-disordered breathing, nocturnal hypoxemia, and subsequent cognitive decline in older community-dwelling men: the osteoporotic fractures in men sleep study. *AmGeriatrSoc* 2015;63:453-461.
2. Bubu OM, Andrade AG, Umasabor-Bubu OQ et al. Obstructive sleep apnea, cognition and Alzheimer's disease: A systematic review integrating three decades of multidisciplinary research. *Sleep Med Rev* 2020 Apr;50:101250.

Literaturverzeichnis

Hamburger Ärzteblatt 12 | 2020

Seite 3

Weidestr. 122 b
22083 Hamburg
Redaktion

E-Mail: verlag@aekhh.de

Tel.: (040) 20 22 99 – 205

Fax: (040) 20 22 99 – 400

3. André C, Rehel S, Kuhn E et al. Association of sleep-disordered breathing with Alzheimer disease biomarkers in community-dwelling older adults: A secondary analysis of a randomized clinical trial. *JAMA Neurol*. Published online March 23, 2020. doi:10.1001/jamaneurol.2020.0311.
4. Liguori C, Mercuri NB, Izzi F et al. Obstructive sleep apnea is associated with early but possibly modifiable Alzheimer's disease biomarkers changes. *Sleep* 2017;40:10.1093.
5. Andrade AG, Bubu OM, Varga AW, Osorio RS. The Relationship between obstructive sleep apnea and Alzheimer's disease. *J Alzheimers Dis* 2018;64 (s1):S255-S270.
6. Richards KC, Gooneratne N, Diccico B et al. CPAP adherence may slow 1-year cognitive decline in older adults with mild cognitive impairment and apnea. *J Am Geriatr Soc* 2019;67:558-564 .
7. Wang Y, Cheng C, Moelter S et al. One year of continuous positive airway pressure adherence improves cognition in older adults with mild apnea and mild cognitive impairment. *Nurs Res* 2020;69:157-164.